eco.mont - Volume 6, Number 2, July 2014 ISSN 2073-106X print version

ISSN 2073-1558 online version: http://epub.oeaw.ac.at/eco.mont

Threats related to tourism and recreation in Natura 2000 areas, based on the analysis of the standard data forms (SDF) from the Alpine Bioregion of Slovakia

Szymon Ciapała, Paweł Adamski, Adam Mroczka, Juraj Svajda & Zbigniew Witkowski

Keywords: Natura 2000 sites, Standard Data Form (SDF), impact of tourism, Slovak Alpine Region

- Abstract -

The aim of this study was to estimate the size and direction of tourism and recreation influence on protected areas within the Natura 2000 sites. The research covered sites in Slovakia within the Alpine Bioregion. Information included in the Standard Data Form (SDF) was analysed and the SDF used as a basic tool to collect data related to the Natura 2000 sites. The tool contains unified information of forms of activities within a given area, their intensity and effect (positive, neutral, or negative) on nature. The study revealed that almost a third of activities pursued in the Slovakian part of the Alpine Bioregion were related to tourism and recreation in the wider sense. Furthermore, the influence of the majority of activities on the environment was considered negative or neutral in all analysed Natura 2000 areas. Only ten forms of activities were found positive in some parts of the Natura 2000 sites. The findings also indicated that the intensity of the analysed activities was similar within the Natura 2000 sites and in the adjacent areas. However, the effects of the selected activities were more often found to be negative within the Natura 2000 sites than in their adjacent areas. The results obtained from this study suggest that broadly understood tourism and recreational activities can be considered an inherent element of human influence on biodiverse areas. They should be taken into consideration in managing and planning Natura 2000 sites and their adjacent areas.

Introduction

Tourism is often perceived as one of the most important elements of sustainable development (Kombol 2000; Krnacova et al. 2001). This form of activity combines the opportunities of economic development with the preservation of high environmental values of the area (Maikihuri et al. 2000; Neto 2003). At the same time, however, an excessive touristic load or its incorrect distribution in the region might result in the degradation of its environmental resources (Hresko & Bugar 2001; Gorczyca & Krzemien 2002).

The most remarkable examples of this phenomenon are the effects of the development of touristic infrastructure in the form of hostels, hotels, roads and railways, ski lifts and ski runs (Witkowski 1996; Łajczak 1996). The impact of tourism on the natural environment has recently been the subject of various (more or less comprehensive) studies (Lidl 1997). This interest stems from the mass character of tourism on the one hand (Freitag & Pyka 2011), and the fact that attractive natural areas have also become popular destinations due to the growing interest in ecotourism, on the other (Reiter 2011). Therefore, among recreational areas, the protected areas, also included in the Natura 2000 ecological network, have begun to play a special role as well. These areas are monitored for the impact of various factors on the condition of the environment. On the basis of inventory studies, the person who submits the area to the Natura 2000 network is asked to fill in a Standard Data Form (SDF) on activities carried out in Natura 2000 areas as well as in the adjacent areas. The SDF has eight main sections, of which the sixth includes data on the possible threats or on activities with positive outcomes for the protected area.

The paper presents the results obtained from the analysis of the data in terms of their impact on the protected areas within the Natura 2000 network. The authors focused on activities related to tourism and recreation in the wider sense.

During the preparation of the Natura 2000 ecological network in Europe, the European Commission outlined the division of the territory into several biogeographic regions (Makomaska-Juchiewicz & Tworek 2003; Dubiel et al. 2010; Wójcik 2010). Our study was limited to the areas of the Alpine Bioregion as defined by the EU nomenclature (ETC/BD 2006).


Methods

The paper is based on information included in the SDF for the sites established in the Slovakian part of the Carpathians (http://www.eea.europa.eu/data-and-maps/data/natura-2000). This source determines the accuracy and type of data presentation. The names and codes of activity types as well as categories of their intensity and effects were based on the EU instructions for the sites delimitation (Natura 2000 SDF Explanatory Note 1999).

The activities related to broadly understood tourism and recreation were selected from instructions in the SDF explanatory notes and further analyses carried out on these activities to establish whether their intensity and effects are similar in the Natura 2000 sites and in adjacent areas. The analyses were conducted by comparing the number of indicators within each intensity category and the direction of each selected activity. Frequency tests were applied to perform comparisons such as Pearson's Chi square, and Likelihood Ratio tests when data quantity did not meet test requirements (Sokal & Rohlf 1995).

Results

Based on the SDFs for all 285 sites in the Slovak part of the Alpine Bioregion, a total of **148** types of human activities with positive and/or negative effect on the Natura 2000 areas were identified in the Alpine Bioregion of a total number of 1397 number activities. Furthermore, **150** types of activities affecting the areas adjacent to the Natura 2000 sites were also de-



■ all types of activities ■ tourism and recreation-related activities

Figure 1 — The number of human activity types inside the

Natura 2000 areas and in the adjacent areas (the Alpine Bioregion).




Figure 2 — The most frequent tourism-related activities reported from the 285 Natura 2000 areas as taking place inside the Natura 2000 sites (blue columns) and in the areas adjacent areas (red columns).




Figure 3 – The activities which most intensively affected the 285 Natura 2000 areas in the Alpine Bioregion (blue columns) and in the areas adjacent to the them (red columns).

termined. However, only a few activities were tourism and recreation-related. After analysing the SDFs, the authors distinguished **45** types of activities inside the Natura 2000 areas and **46** types of activities in the adjacent areas (Figure 1).

The number and forms of tourism and recreation-related activities

Almost one third of all 150 types of activities carried out in the Alpine areas were associated with tourism. Not only the types of tourism-related activities were taken into consideration but also the activities which enable and facilitate the realization of touristic and recreational needs.

Table 1 shows the intensity of various types of tourism and recreation-related activities within the Natura 2000 areas in the Alpine Bioregion. The intensity of touristic and recreational activities and their effects were presented on a 3-point scale, in line with the SDF-adopted principles (Natura 2000 SDF Explanatory Note 1999).

Threats to the Natura 2000 environmental values may not only result from the tourism and recreation-related activities pursued by people within the boundaries of a particular area, but also from activities carried out outside it in the adjacent areas. Table 2 shows the sources of such threats. However, in relation to the Table 1, a comparable variety of tourism and recreation-related activities which might pose a threat can be observed, and two other types of activities which do not take place inside the Natura 2000 area. These include position 16 (Airfields and helipads) and 23 (Stadiums) in Table 2. Given their rarity, these activities seem insignificant.

Data analysis

The analysis of the information obtained from the SDFs indicated that the particular threats associated with tourism and recreation in the majority of areas occur within their boundaries as well as in their adjacent areas.

The results obtained from the SDFs indicated that particular types of activities can vary in terms of the intensity of their impact on the area, see Figure 3.

The majority of human activities had a negative and only two (educational centres and tunnels) a positive effect, primarily on the areas in which they were located. Only 10 types of activities were considered positive in 28 cases of completed SDFs. They represent less than 2% of all (1 397 cases) touristic and recreational activities in the Alpine Bioregion.

The analysis of the intensity of activities inside and in the vicinity of the Natura 2000 sites returned significant differences for only two types of activities: vandalism (code 740), with significantly higher intensity inside the site than in the vicinity (p = 0.0125), and, in contrast, invasion of alien species (code 954), with significantly weaker or average intensity inside the site (p = 0.0062).

Table 1 – Number of the Natura 2000 areas where tourism and recreation-related activities were recorded (the Alpine Bioregion). Source: authors' own studies

No.	Activity	Activity type	Intensity	of activit	у	Effects of activity		
			strong	average low		positive neutral		negative
	code		(A)	(B)	(C)	(+)	(0)	(-)
1	220	Fishing	10	21	18	2	27	20
2	230	Hunting	37	76	42	2	104	49
3	240	Taking and removal of animals in general	3	4	7	3	5	6
4	241	Taking and removal insects, amphibians, or reptiles	2	7	26	0	4	31
5	250	Plant collection in general	1	10	3	0	4	10
,	051	Collection of perennials, forest fruits and	1.5	E 7	48	0	7.4	44
6	251	medicinal plants	15	57	40	0	76	44
7	7 290	Other forms of hunting, fishing and collect-	0	6	6	3	3	6
		ing specimens of animals or plants						
8	403	Scattered housing	0	3	15	5	12	1
9	420	Waste, sewage	0	3	5	0	3	5
10	421	Waste and municipal sewage	1	16	13	0	4	26
11	500	Transport network	2	20	5	0	4	23
12	501	Paths, hiking trails, bike trails	7	39	50	0	59	37
13	502	Roads, highways	3	20	17	0	13	27
14	503	Railways	1	4	8	0	8	5
15	506	Airfields, helipads	0	0	3	0	1	2
16	507	Bridges and viaducts	0	4	13	0	10	7
17	508	Tunnels	1	0	4	3	2	0
18	509	Other transport networks	3	9	5	0	5	12
19	600	Sports and recreational infrastructure	1	10	11	0	9	13
20	601	Golf courses	0	0	1	0	0	1
21	602	Ski resorts	1	10	13	0	11	13
22	604	Running and racing tracks	0	0	1	0	0	1
23	605	Equestrian areas	0	1	0	0	0	0
24	606	Leisure parks	1	1	0	0	1	1
25	607	Sports fields	0	1	7	0	6	2
26	608	Campsites and caravan parks	0	3	5	0	3	5
27	609	Other sports and recreational areas	0	6	9	0	7	8
28	610	Educational centres	1	1	4	4	2	0
29	620	Outdoor sports and activities	4	12	12	0	9	19
30	621	Water sports	5	4	6	0	5	10
31	622	Hiking, horse riding and biking tourism	14	58	59	0	78	53
32	623	Off-road and motorcycles	6	19	21	0	2	44
33	624	Mountaineering, rock climbing, caving	7	24	18	0	21	28
34	625		4	14	11	0	5	24
35	626		2	12	10	0	8	16
37	629	Other sports and recreational activities in nature	2	9	10	0	7	14
38	690	Other possible effects of recreational and sport activities	1	3	3	0	5	2
39	720		7	20	7	0	1	33
40	740		8	18	11	0	2	35
41	870	Dams, embankments, use of river banks	5	7	9	4	5	12
42	900		12	39	18	1	11	57
43	942	Avalanches and other mass movements	3	6	3	1	9	2
44	952		5	15	13	0	4	29
45		Invasion of alien species	5	39	36	0		71

In terms of activity effects, almost 20% of selected activity types showed significant differences. In 8 types of activities: collection of perennials, forest fruits and medicinal plants (p = 0.0481); transport network (p = 0.0100); paths, hiking trails, bike trails (p = 0.0080); other transport networks (p = 0.0248); water sports (p = 0.0284); hiking, horse riding and biking tourism

(p=0,0010); other sports and recreational activities in nature (p=0,0107); dams, embankments, use of river banks (p=0,0449). Activities inside the Natura 2000 sites more often referred to as negative than in their vicinity. However, tunnels (code 508) were perceived as having a significantly more positive effect inside the Natura 2000 sites than in their vicinity (p=0,0297).

Table 2 – Number of areas adjacent to Natura 2000 sites where tourism and recreation-related activities were recorded (the Alpine Bioregion) Source: authors' own studies.

No.	Activity code	Activity type	Intensity of activity			Effects of activity		
			strong (A)	average (B)	low (C)	positive (+)	neutral (0)	negative (-)
1	220	Fishing	5	30	17	1	44	7
2	230		39	74	44	4	113	40
3	240	Taking and removal animals in general	1	4	6	1	1	9
4	241	Taking and removal insects, amphibians, or reptiles	0	3	11	0	2	12
5	250	Plant collection in general	1	10	3	0	4	10
6	251	Collection of perennials, forest fruits and medicinal plants	15	44	40	1	75	23
7	290	Other forms of hunting, fishing and collecting specimens of animals or plants	1	4	4	0	4	5
8	403	Scattered housing	0	7	9	1	15	0
9	420	Waste, sewage	4	12	6	0	4	18
10	421	Waste and municipal sewage	6	34	15	0	7	48
11	500	Transport network	10	17	9	0	16	20
12	501	Paths, hiking trails, bike trails	1	28	38	0	54	13
13	502	Roads, highways	24	63	32	0	42	77
14	503	Railways	11	21	15	0	24	23
15	505	Airports and airport infrastructure	0	3	1	0	1	3
16	506	Airfields, helipads	0	0	10	0	3	7
17	507	Bridges and viaducts	0	9	7	0	8	8
18	508	Tunnels	1	0	3	0	4	0
19	509	Other transport networks	1	5	5	0	8	3
20	600		7	20	12	0	20	19
21	601	Golf courses	0	2	0	0	2	0
22	602	Ski resorts	5	12	6	0	9	14
23	603		1	1	4	0	4	2
24	604		1	0	1	0	0	2
25	605	Equestrian areas	1	2	1	0	3	1
26	606		1	0	1	0	0	2
27	607		1	5	11	0	14	3
28	608	'	2	7	13	0	11	11
29	609		3	9	10	0	10	12
30	610	Educational centres	1	0	7	6	1	1
31	620	Outdoor sports and activities	8	21	9	0	22	16
32	621	Water sports	3	3	6	0	9	3
33	622	Hiking, horse riding and biking tourism	9	33	61	0	82	21
34	623		7	20	16	0	4	39
35		Mountaineering, rock climbing, caving	4	7	8	0	10	9
36	625		5	6	12	0	8	15
37	626	Skiing, including off-piste, ski touring	4	9	11	0	12	12
38	629	Other sports and recreational activities in nature	3	8	8	0	14	5
39	690	Other possible effects of recreational and sport activities	1	2	2	0	3	2
40	720	Trampling; excessive exploration	3	9	2	1	0	13
41	740	Vandalism	0	15	9	0	3	21
42	870	Dams, embankments, use of river banks	6	10	4	0	4	16
43	900		6	24	15	1	8	36
44	942	Avalanches and other mass movements	0	1	1	0	1	1
45	952		10	17	9	0	2	34
46	954		22	44	30	0	7	89

In terms of activity effects, almost 20% of selected activity types showed significant differences. In the majority of the cases, activities inside the Natura 2000 sites more often referred to as negative than in their

vicinity. However, tunnels (code 508) were perceived as having a significantly more positive effect inside the Natura 2000 sites than in their vicinity.

Discussion

Of 45 types of activities mentioned inside or in the vicinity of the Natura 2000 sites, only 10 were considered positive in 28 cases of completed SDFs. The most significant types of human activities, related to a broadly understood tourism and recreation, and having a positive effect on the Natura 2000 areas were: the existence of scattered housing – 5 cases; construction of dams, shafts and use of river banks – 4 cases, and the impact of educational centres located in the Natura 2000 areas – 4 cases. In the adjacent areas to the Natura 2000 sites, positive types of tourism and recreation-related activities were activities of educational facilities – 6 cases, and hunting – 4 cases.

The results show that the high impact intensity of a given factor may not necessarily be a basis for concern or for the conclusion that it poses a threat to the area. High intensity factors, although often associated with a negative influence on the environment, may not necessarily affect it negatively every time. For instance – hunting, which is the most frequently mentioned factor affecting the Natura 2000 areas and the adjacent areas, and the most intensive type of activity A (strong): in 1% of cases is it treated as having a positive effect, in 67% as neutral, and only in 32% of all areas is it perceived as having a negative effect.

Hunting represents 1.3% of all 28 cases of positive effects, 18% of all 565 activities are seen as neutral and 6.1% of all 804 as having negative effects. In spite of the fact that hunting is mentioned as the most frequent activity and usually with a strong influence on the area, in the light of the data obtained from SDFs it does not pose any threat to the protected Natura 2000 areas.

The comparison between Natura 2000 and adjacent areas shows that generally there are no differences in intensity for the activity types analysed. The significant differences in terms of intensity were noted only for invasive alien species and vandalism. The higher intensity of threats related to invasive alien species inside the Natura 2000 results from the absence of a protective regime that would prevent alien species from growing and spreading, and a higher density of settlements accompanied by green infrastructure outside the Natura 2000 sites. Research indicates that settlements are regarded as the main source of alien species invasions to protected areas (McNeel 2001). However, the reason for higher intensity of vandalism inside the Natura 2000 is less known. There is little evidence that behaviour considered as vandalism happens more often inside the sites than in their vicinity. The difference may be explained by the fact that such behaviour, e.g. picking up flowering plants inside the Natura 2000 sites, are not perceived as vandalism in the adjacent areas.

While no significant differences in the analysis of activity intensity inside the Natura 2002 sites and in the adjacent areas were observed, the discrepancy in

the analysis of activity effects was noted. It may be explained in two ways. First, it is assumed that the same intensity of a given factor causes a stronger environmental reaction inside the Natura 2000 sites, e.g. bigger participation of primal and more sensitive ecosystems within the site that in the adjacent site. Second, the discrepancy may be caused by the practice of filling out the SDFs. Instructions related to the assessment of intensity and effects of activities are not entirely objective. Thus, a factor of the same intensity may be assessed differently in terms of its effect.

Both mechanisms are perhaps responsible for indicating negative effects of the same forms of activities more often inside than outside the Natura 2000 sites. The only factor indicated as positive more frequently within the Natura 2000 site than outside it, are tunnels (code 508). This difference stems from the fact that tunnels not only are constructed due to technical requirements of road development, but also to enable animal migration.

The results obtained from this study indicate that broadly understood tourism and recreational activities are an inherent element of human influence on biodiverse areas. Therefore, they should be taken into consideration when managing the Natura 2000 sites and for their adjacent areas.

References

Dubel, A., M. Jamontt-Skotis, K. Królikowska, J. Stefańska & A. Banrowska 2010. *Skuteczne zarządzanie obszarami Natura 2000*. Stowarzyszenie Centrum Rozwiązań Systemowych, Wrocław-Kraków: 37–49.

ETC/BD – European Topic Centre on Biological Diversity 2006. *The indicative Map of European Biogeographical Regions: Methodology and development.* Muséum National d'Histoire Naturelle. Paris.

Freitag, R. & D. Pyka 2011. Status Quo and Forecasting World and European Tourism. In: Roland, C. & M. Buck (eds.), *Trends and Issues in Global Tourism*: 3–14

Gorczyca, E. & K. Krzemień 2002. Wpływ ruchu turystycznego na rzeźbę Tatrzańskiego Parku Narodowego. In: Borowiec, W., A. Kotarba, A. Kownacki, Z. Krzan & Z. Mirek (eds.), Przemiany środowiska przyrodniczego Tatr. Tatrzański Park Narodowy, Polskie Towarzystwo Przyjaciół Nauk o Ziemi, Oddział Kraków, Kraków – Zakopane: 389 –394.

Hresko, J. & G. Bugar 2001. Problems of natural hazard assessment and monitoring in the Tatra Mts. *Ekologia-Bratislava* 20: 96–100.

Kombol, P. 2000. Rural tourism on the Croatian islands – Sustainable development and regenerative strategies. *Periodicum Biologorum* 102: 425–431.

Krnacova, Z, K. Pavlickova & P. Spisiak 2001. The assumptions for the tourism development in rural areas in Slovakia. *Ekologia-Bratislava* 20 (3): 317–324.

Lidle M. 1997. Recreation ecology. London.

Lajczak A. 1996. Wpływ narciarstwa i turystyki pieszej na erozję gleby w obszarze podszczytowym Pilska. In: Łajczak, A., S. Michalik & Z. Witkowski (eds.), Wpływ narciarstwa i turystyki pieszej na przyrodę Masywu Pilska. Studia Naturae 41: 131–159.

Makomaska-Juchiewicz, M. & S. Tworek 2003. Miejsce sieci Natura 2000 w europejskiej ochronie przyrody. In: Makomaska-Juchiewicz, M. & S. Tworek (eds.), Ekologiczna sieć Natura 2000, problem czy szansa. Instytut Ochrony Przyrody PAN, Kraków: 9–21.

Maikhuri, R.K., U. Rana, K.S. Rao, S. Nautiyal & K.G. Saxena 2000. Promoting ecotourism in the buffer zone areas of Nanda Devi Biosphere Reserve: an option to resolve people-policy conflict. *International Journal of Sustainable Development and World Ecology* 7 (4): 333–342.

McNeel, J.A. 2001. An Introduction to Human Dimensions of Invasive Alien Species. In: McNeel, J.A. (ed.), *The Great Reshuffling Human Dimensions of Invasive Alien Species*. IUCN. Cambridge: 4–20.

NATURA 2000 Standard Data Form Explanatory Notes 1999. Available at: http://ec.europa.eu/environment/nature/legislation/habitatsdirective/docs/standarddataforms/notes_en.pdf (accassed 01/03/2014)

Neto, F. 2003. A new approach to sustainable tourism development: Moving beyond environmental protection. *Natural Research Forum* 27 (3): 212–222.

Reiter, A. 2011. Eco-leadership and Green Lifestyle: Successful Strategy for a Growing Market Segment? In: Roland, C. & M. Buck (eds.), *Trends and Issues in Global Tourism*: 93–96.

Sokal, R.R. & J.R. Rohlf 1995. Biometry: The Principles and Practices of Statistics in Biological Research. Third Edition. New York.

Witkowski, Z. 1996. Narciarstwo i ochrona przyrody na Pilsku: tlo i przyczyny konfliktu. In: Łajczak, A., S. Michalik & Z. Witkowski (eds.), *Wpływ narciarstwa i turystyki pieszej na przyrodę Masywu Pilska*. Studia Naturae 41: 5–12.

Wójcik, B. 2010. Natura 2000 – podstawowe informacje. In: Kamieniecka, J. & B. Wójcik (eds.), *Natura 2000. ABC dla turystyki*. Instytut na rzecz Ekorozwoju, Warszawa: 13–14.

Authors

Szymon Ciapała

Head of Ecology Faculty; Academy of Physical Education in Kraków. His scientific interests focus on the issues of tourist impact on nature and methods of tree ring dating as assessment of the touristic impact.

Paweł Adamski

Researcher at the Institute of Nature Conservation PAS. Specializes in the implementation of the population and behavioural ecology into nature conservation practices, also interested in environmental impact of the tourism and recreation.

Adam Mroczka

Lecturer at the Ecology Faculty, Academy of Physical Education in Cracow. His research interests focus on environmental conditions of tourist movements, the influence of the environment on human health and bioclimatic conditions of human settlements.

Juraj Svajda

Assistant at Matej Bel University in Banská Bystrica (Slovakia). His research interests focus on Protected Area Management including monitoring of visitors in national parks.

Zbigniew Witkowski

Head of Department of Environmental Sciences. Author of over 270 publications on nature conservation and tourism, interested in the management of tourism in the context of nature conservation in protected areas.